


## *Streptomyces* sp. CBB-KSK145

|  |  |  |
|--|--|--|
| Colony morphology                        | Greyish white aerial mycelium; circular; entire; umbonate; dry; opaque; medium colony size<br> |  |
| Gram-stain                               | Positive, filamentous bacterium  |  |
| Isolation medium                         | Starch Casein Agar (SCA) + 50% Artificial sea water (ASW)  |  |
| Growth condition                         | Aerobic; pH7; 28±2°C; 7-14 days  |  |
| Sampling date, location, source          | 12 May 2014; Kuala Sala, Kedah (100.3546°E, 5.9737°N); Sediment  |  |
| Biochemistry/Physiology                  | No   |  |
| 16s rRNA gene analysis(EzBioCloud/N CBI) | Top-hit taxon  | <i>Streptomyces pactum</i>   |
|  | Top-hit strain   | NBRC 13433(T)  |
|  | Similarity (%)   | 99.92  |
|  | Top-hit taxonomy   | Bacteria;Actinobacteria;Actinobacteria_c;Streptomycetales;Streptomycetaceae;Streptomyces |
| Publication                              | No   |  |
| Application                              | No   |  |
| Isolated by                              | Farizah Hanim bt Lat   |  |
| Risk group                               | 2  |  |
| Additional information                   |  |  |